Author Index

AARDEN L., CORLEY R.B., SÖDERBERG A. & LEFKOVITS I. Limiting dilution analysis of the suppressive effect mediated by alloantigen-primed cells, 399
AARDEN L.A., *see* LEFKOVITS I.
ABATÁNGELO CARMEN, *see* MARGNI R.A.
AHMED A., *see* HACE J.A.
ANTOINE J.-C, *see* GOUD B.
ASHBY J., *see* EREMIN O.
AVNI A., *see* SHOHAM J.
AVRAMEAS S., *see* GOUD B.

BAHR G.M., ROOK G.A.W., MORENO E., LYDYARD P.M., MODABBER F.Z. & STANFORD J.L. Use of the ELISA to screen for anti-thymocyte and anti-β_2 microglobulin antibodies in leprosy and SLE, 865
BALDWIN R.W., *see* BROOKS C.G.
BALTZ MARILYN L., DYCK R.F. & PEPYS M.B. Amyloid P-component in mice injected with casein: identification in amyloid deposits and in the cytoplasm of hepatocytes, 59
BASCH R.S., *see* BUMOL T.F.
BATCHELOR K.W. & STANWORTH D.R. A comparison of the histamine-releasing properties of rat pleural and peritoneal mast cells, 271
BATTY J.E., *see* HOLT P.G.
BAX J., *see* BERG W.B. VAN DEN
BEARD, HELEN K. RYVAR RUTH, BROWN, R. AND MUIR HELEN. Immunochemical localization of collagen types and proteoglycan in pig intervertebral discs, 491
BEFUS A.D., *see* DENBURG J.A.
BEFUS A.D., *see* BIENENSTOCK J.
BENNER R., *see* HAAIJMAN J.J.
BENOLIEL ANNE MARIE, CAPO C., BONGRAND P., RYTER ANTOINETTE & DEPIEDS R. Non-specific binding by macrophages: existence of different adhesive mechanisms and modulation by metabolic inhibitors, 547
BERG W.B. VAN DEN, HAASAKKER TINEKE C., BAX J. & SCHEPER R.J. Suppression of antibody-mediated accumulation of eosinophils in chronic inflammatory lesions by concomitant delayed hypersensitivity reactions, 981
BERTOVICH M.J., *see* MACDERMOTT R.P.
BHAKDI S. & TRANUM-JENSEN J. Re-incorporation of the terminal C56-9 complement complex into lipid bilayers: formation and stability of reconstituted liposomes, 737
BIENENSTOCK J., *see* DENBURG J.A.
BIENENSTOCK J. & BEFUS A.D. Mucosal immunology, 249
BINAGHI R.A., *see* MARGNI R.A.
BIRMINGHAM J.R. & JESKA E.L. The isolation, long-term cultivation and characterization of bovine peripheral blood monocytes, 807
BITO Y., *see* SAWADA T.
BONGRAND P., *see* BENOLIEL ANNE MARIE
BOURNE F.J., *see* MORGAN K.L.
BOURNE F.J., *see* NEWBY T.J.
BRAIDWOOD E.A., *see* HENDERSON D.C.
BRITTLE M.P. & PLAYFAIR J.H.L. The role of T cells in regulating immunoglobulin class and subclass, 743
BROOKS C.G., FLANNERY G.R., WEBB P.J. & BALDWIN R.W. Quantitative studies of natural immunity to solid tumours in rats. The nature of the killer cell depends on the type of assay, 673
BROOM B.C. & SOUTHERN M.A. Evidence for control of immune cell arrival at sites of antigen challenge in previously immunised animals, 523
BROWN R., *see* BEARD H.
BRUCE R.G., *see* OTTAWAY C.A.
BRULEY-ROSSET M., *see* KIGER N.
BUMOL T.F., RETZEL E.F., DOUGLAS S.D., BASCH R.S., BUXBAUM J.N. & FARAS A.J. Terminal deoxynucleotidal transferase expression in a Thy-1 alloantigen variant lymphoma cell line, 799
BUTCHER G.W., *see* HOWARD J.C.
BUXBAUM J.N., *see* BUMOL T.F.

CALDWELL J., *see* LEANER T.
CAPO C., *see* BENOLIEL ANNE MARIE
CATOVSKY D., *see* FOA R.
CHALLACOMBE S.J., *see* LEHNER T.
CHANDALI YARDENA, *see* SHOHAM J.
CHATURVEDI U.C., DALAKOTI HEMLATA & MATHUR ASHA Characterization of the cytotoxic factor produced in the spleen of dengue virus-infected mice, 387
CHIN D.K., *see* KIM B.S.
CLARKSON CAROL A., *see* SYMONS D.B.A.
COHEN MIRIAM, *see* SHOHAM J.
COLE P., *see* SAINT-REMY J.-M.R.
COLE P.J., *see* EASMON C.S.F.
COLE P.J., *see* WILLIAMS A.J.
COLLET H., *see* GAUSSET P.H.
COLTEN H.R., *see* PASSWELL J.H.
COOKE ANNE & HUTCHINGS PATRICIA. Sex differences in the regulation of experimentally induced autoantibodies in (NZB × NZW)F_1 mice, 819
COOKE ANNE, HUTCHINGS PATRICIA & MARSHALL-CLARKE S. Lack of auto antigen-specific splenic suppressor cells in mice with an X-linked B-lymphocyte defect, 815
COOMBS R.R.A., *see* EREMIN O.
CORLEY R.B., *see* LEFKOVITS I.
CORLEY R.B., *see* AARDEN L.
CORTENS M., SKLAR SUZI & RICHTER M. The detection and identification of subpopulations of circulating human lymphocytes, monocytes and neutrophils capable of effecting a mitogen-induced cell-mediated cytotoxic reaction towards erythrocytes of various species, 623

DAHA M.R., *see* KIJLSTRA A.
DAHA M.R. & ES L.A. VAN. Isolation and characterization of

rat complement factor B and its interaction with cell bound human C3, 849
DALAKOTI HEMLATA, *see* CHATURVEDI U.C.
DE BAETSELIER P., VAECK M., DE SMET W. & RON Y. Antigen-induced proliferation assay for rabbit T lymphocytes. I. Characteristics of the response, 997
DE CEULAER C., PAPAZOGLOU S. & WHALEY K. Increased biosynthesis of complement components by cultured monocytes, synovial fluid macrophages and synovial membrane cells from patients with rheumatoid arthritis, 37
DE SMET W., *see* DE BAETSELIER P.
DEL PRETE G.F., *see* ROMAGNANI S.
DELESPESSE G., *see* GAUSSET PH.
DENBURG J.A., BEFUS A.D. & BIENENSTOCK J. Growth and differentiation *in vitro* of mast cells from mesenteric lymph nodes of *Nippostrongylus brasiliensis*-infected rats, 195
DEPIEDS R., *see* BENOLIEL ANNE MARIE
DEVEY MADELEINE E. & STEWARD M.W. The induction of chronic antigen–antibody complex disease in selectively bred mice producing either high or low affinity antibody to protein antigens, 303
DIAMANTSTEIN T., TRISSL DOROTHEA, KLOS M., GOLD D. & HAHN H. Mitogenicity of *Entamoeba histolytica* extracts for murine lymphocytes, 347
DOCKRELL HAZEL M., SOUZA J.B. DE & PLAYFAIR J.H.L. The role of the liver in immunity to blood-stage murine malaria, 421
DOLBY JEAN M., *see* STEPHENS SUSAN
DONATSCH P., *see* RYFFEL B.
DORP W. VAN, *see* KIJLSTRA A.
DOUGLAS S.D., *see* BUMOL T.F.
DRESSER D.W. & POPHAM ANGELA M. Rheumatoid factor in mice: plaque assay for homophile and heterophile rheumatoid factors, 569
DRESSER D.W., *see* POPHAM ANGELA M.
DUCHATEAU J., *see* GAUSSET PH.
DYCK R.F., *see* BALTZ MARILYN L.

EASMON C.S.F., *see* WILLIAMS A.J.
EASMON C.S.F., COLE P.J., WILLIAMS A.J. & HASTINGS M. Measurement of opsonic and phagocytic function by Luminol-dependent chemiluminescence, 67
ENGERT R.F., *see* KLIPSTEIN F.A.
EREMIN O., COOMBS R.R.A., ASHBY J. & PLUMB D. Natural cytotoxicity in the guinea-pig: the natural killer (NK) cell activity of the Kurloff cell, 367
ES L.A. VAN, *see* DAHA M.R.

FALAGIANI P., *see* ROMAGNANI S.
FARAS A.J., *see* BUMOL T.F.
FLANNERY G.R., *see* BROOKS C.G.
FOA R., ZAFAR M.N. & CATOVSKY D. Inhibition of T-lymphocyte colony formation by inhibitors of mitochondrial protein synthesis, 55
FOLKERD ELIZABETH J., GARDNER BRIGITTE & HUGHES-JONES N.C. The relationship between the binding ability and the rate of activation of the complement component C1, 179
FORMEISTER J.F., *see* MACDERMOTT R.P.
FEDERIX-WOLTERS ELIZABETH M.H., *see* RIJKERS G.T.
FREITAS A.A., *see* ROCHA BENEDITA

GALFRÈ G., *see* HOWARD J.C.
GARDNER BRIGITTE, *see* FOLKERD ELIZABETH J.
GAUSSET PH., DELESPESSE G., DUCHATEAU J. & COLLET H. *In vitro* response of human peripheral blood lymphocytes to protein A DNA synthesis and generation of cells synthesizing the three major classes of Ig, 891
GENTILE TERESA, *see* MARGNI R.A.
GERDES J. & STEIN H. Physicochemical characterization of C3b receptors isolated from human erythrocytes by immunoprecipitation, 929
GILMORE R.J., *see* RENSHAW H.W.
GOETZL E.J., *see* VALONE F.H.
GOLD D., *see* DIAMANTSTEIN T.
GONATAS N.K., *see* GOUD B.
GONDA N., *see* KANO S.
GÖTZ U., *see* RYFFEL B.
GOUD B., ANTOINE J.-C., GONATAS N.K., STIEBER ANNA & AVRAMEAS S. A morphological and functional study on antigen binding and endocytosis by immunocytes, 833
GUTIÉRREZ CARMEN., *see* MARTIN MARIA J.
GYURE L.A., *see* HALL J.G.

HAAIJMAN J.J., SLINGERLAND-TEUNISSEN JOKE, OUDENAREN A. VAN, MINK J.G. & BENNER R. Kinetics of recovery of serum Ig levels and of cytoplasmic Ig positive cells in various lymphoid organs of nude mice after thymus transplantation, 279
HAASAKKER TINEKE C., *see* BERG W.B. VAN DEN
HAEGERT D.G. Demonstration of immunoglobulin-related molecules on human peripheral blood T lymphocytes using chicken anti-F(ab′)$_2$ and anti-IgM antibodies, 663
HAHN H., *see* DIAMANTSTEIN T.
HALL J.G., GYURE L.A. & PAYNE A.W.R. Comparative aspects of the transport of immunoglobulin A from blood to bile, 899
HARE J.A., AHMED A. & SELL K.W. *In vitro* and *in vivo* response of lymphoid cells from LHC hamsters to murine thymus-independent and thymus-dependent antigens, 705
HASTINGS M., *see* EASMON C.S.F.
HASTINGS M.J.G., *see* WILLIAMS A.J.
HENDERSON D.C., BRAIDWOOD E.A., SMILLIE D.M. & WHITE R.G. Kinetics of the decline phase of the primary response to human serum albumin in the bird, 449
HERLES SUSAN, *see* SCHONFELD S.E.
HILL H.R., *see* OGDEN B.E.
HIMENO K., *see* NOMOTO K.
HOGG NANCY & PARISH C.R. Surface antigens of the murine cytostatic peritoneal macrophage, 187
HOLT P.G. & BATTY J.E. Alveolar macrophages. V. Comparative studies on the antigen presentation activity of guinea-pig and rat alveolar macrophages, 361
HORMAECHE C.E. The *in vivo* division and death rates of *Salmonella typhimurium* in the spleens of naturally resistant and susceptible mice measured by the superinfecting phage technique of Meynell, 973
HOWARD C.J. Variation in the susceptibility of bovine mycoplasmas to killing by the alternative complement pathway in bovine serum, 561
HOWARD J.C., BUTCHER G.W., LICENCE DIANA R., GALFRÈ G., WRIGHT B. & MILSTEIN C. Isolation of six monoclonal alloantibodies against rat histocompatibility antigens: clonal competition, 131
HUCHET R., *see* KIGER N.

HUGHES-JONES N.C., *see* FOLKERD ELIZABETH J.
HUSSEIN A.M., *see* MORGAN K.L.
HUTCHINGS PATRICIA, *see* COOKE ANNE

JACOBSEN C., *see* JOHANSEN A.S.
JESKA E.L., *see* BIRMINGHAM J.R.
JOHANSEN A.S., STEENSGAARD J. & JACOBSEN C. Solubility properties of IgG immune complexes. Comparison between the effects of low molecular weight solvents and polyethylene glycols, 695
JOSHI D.S., *see* SELVAN EMILY

KAKIUCHI T., *see* MIZUGUCHI J.
KANO S., OSHIMI K., SUMIYA M. & GONDA N. Activation of secondary cytotoxic lymphocytes by cell-free factors from *I*-region-primed and *D*-region-primed lymphocytes, 653
KATZ D.H. Recent studies on the regulation of IgE antibody synthesis in experimental animals and man, 1
KAUSHIK A.K. & PANDEY R. Cellular and humoral immune responses to buffalopox virus in experimentally infected mice and rabbits, 153
KIGER N., BRULEY-ROSSET M., HUCHET R. & MATHE G. Study of the effect of single or multiple doses of BCG on antibody-dependent cellular cytotoxicity, 687
KIJLSTRA A., VAN DORP W., DAHA M.R. & LESLIE R.G.Q. The effect of lidocaine on the processing of soluble immune aggregates and immune complexes by peritoneal macrophages, 237
KIM B.S. & CHIN D.K. Presence of a tumour-inhibiting factor (TIF) in sera from normal but not tumour-bearing mice, 289
KITAMURA H., *see* NAGAKI K.
KLIPSTEIN F.A., ENGERT R.F. & SHORT HELEN B. Immunological cross-reactivity of heat-labile enterotoxins produced by enterotoxigenic and enteropathogenic strains of *Escherichia coli*, 115
KLOS M., *see* DIAMANTSTEIN T.
KNOP J. Influence of various macrophage populations on Con-A induced T-cell proliferation, 379
KONIG W., *see* MOLLER G.
KONTIAINEN SIRKKA, *see* ZANDERS E.D.
KUBO C., *see* SHIMAMOTO Y.
KUROIWA A., *see* SAKEMI T.

LACHMANN P.J., OLDROYD R.G., MILSTEIN C. & WRIGHT B.W. Three rat monoclonal antibodies to human C3, 503
LAMB J.R., *see* ZANDERS E.D.
LAMBERT FRANCINE, *see* LIACOPOULOS MONIQUE
LAVROVSKY V.A. & VIKSLER V. KH. Natural killer activity of Wistar rat spleen cells: blocking effect of homologous sera, 541
LEE C.J. Maternal–foetal interaction, antibody formation, and metabolic response in mice immunized with pheumococcal polysaccharides, 45
LEFFORD M.J. Properties of peritoneal exudate lymphocytes that mediate tuberculin delayed-type hypersensitivity and anti-tuberculosis immunity. II. Velocity sedimentation analysis, 643
LEFFORD M.J. Properties of peritoneal exudate lymphocytes that mediate tuberculin delayed-type hypersensitivity and anti-tuberculosis immunity. I. The effect of cytotoxic agents, 635
LEFKOVITS I., *see* AARDEN L.
LEFKOVITS I., AARDEN L. & CORLEY R.B. Ratio-dominance model of suppression: an analysis by limiting dilution, 407
LEHNER T., *see* SCULLY C.M.
LEHNER T., CHALLACOMBE S.J. & CALDWELL J. Oral immunization with *Streptococcus mutans* in rhesus monkeys and the development of immune responses and dental caries, 857
LEHNER T., *see* ZANDERS E.D.
LESLIE R.G.Q., *see* KIJLSTRA A.
LIACOPOULOS P., *see* LIACOPOULOS MONIQUE
LIACOPOULOS MONIQUE, LAMBERT FRANCINE & LIACOPOULOS P. Non-specific inhibitory processes of immunological and mitogenic cellular responses. I. Comparative effect of four suppressive agents, 143
LICENCE DIANA R., *see* HOWARD J.C.
LIFSHITZ RUTH, SCHWARTZ M. & MOZES EDNA Specificity of genes contrilling immune responsiveness to (T,G)-A–L and (Phe,G)-A–L, 339
LIND INGA, *see* RHODES JOAN M.
LUKIĆ M.L. & SIMIĆ M.M. Accessory cell function in immune responses *in vivo*: enhancing effect of radioresistant spleen cells on antibody response to SRBC in rats, 323
LYDYARD P.M., *see* BAHR G.M.

MACDERMOTT R.P., NASH G.S., MERKLE NANCY S., WEINRIEB I.J., BERTOVICH M.J. & FORMEISTER J.F. Further evidence that antibody-dependent and spontaneous cell-mediated cytotoxicity are mediated by different processes of cell types, 439
MAGGI E., *see* ROMAGNANI S.
MANSON-SMITH D.F., *see* OTTAWAY C.A.
MARCINKIEWICZ J. & PTAK W. Macrophage suppressor factor in contact sensitivity. Mechanisms of its release and action, 211
MARGNI R.A., PERDIGÓN GABRIELA, ABATÁNGELO CARMEN, GENTILE TERESA & BINAGHI R.A. Immunobiological behaviour of rabbit precipitating and non-precipitating (co-precipitating) antibodies, 681
MAROM Z., *see* PASSWELL J.H.
MARSHALL-CLARKE S., *see* COOKE ANNE
MARTIN MARIA J., VEGA J. & GUTIÉRREZ CARMEN. Antibody-independent complement activation by a subpopulation of mitogen-induced human T lymphoblasts, 535
MATHE G., *see* KIGER N.
MATHUR ASHA, *see* CHATURVEDI U.C.
MATSUMOTO MISAKO, *see* NAGAKI K.
MATUHASI T., *see* MIZUGUCHI J.
MERKLE NANCY S., *see* MACDERMOTT R.P.
MERLER E., *see* PASSWELL J.H.
MILSTEIN G., *see* LACHMANN P.J.
MILSTEIN C., *see* HOWARD J.C.
MILTON J.D. The effect of glutaraldehyde fixation on the immunogenicity of allogeneic lymphoid and tumour cells, 715
MINK J.G., *see* HAAIJMAN J.J.
MITSUHASHI S., *see* SAITO KAZUKO
MIZUGUCHI J., KAKIUCHI T., NARIUCHI H. & MATUHASI T. Effect of lipopolysaccharide (LPS) on the thymocyte response to PHA: strain difference, 393
MODABBER F.Z., *see* BAHR G.M.
MOLLER G. & KONIG W. Binding characteristics of aggre-

gated IgGa to rat basophilic leukaemia (RBL) cells and rat mast cells, 605
MONTREUIL J., *see* STEPHENS SUSAN
MOORE G.E., *see* VESOLE D.H.
MORENO E., *see* BAHR G.M.
MORGAN K.L., HUSSEIN A.M., NEWBY T.J. & BOURNE E.J. Quantitation and origin of the immunoglobulins in porcine respiratory tract secretions, 729
MORRIS D., *see* STEVENSON F.K.
MOZES EDNA, *see* LIFSHITZ RUTH
MUIR H., *see* BEARD H.
MUISWINKEL W.B. VAN, *see* RIJKERS G.T.

NAGAKI K., MATSUMOTO MISAKO & KITAMURA H. A low molecular weight inhibitor of the alternative complement pathway. I. Its isolation from human urine and the reaction mechanism, 789
NAKAMURA K. Conversion of immune response patterns from high to low and low to high by an RNase-sensitive thymocyte extract, 25
NARIUCHI H., *see* MIZUGUCHI J.
NASH G.S., *see* MACDERMOTT R.P.
NATSUNO T., *see* SAITO KAZUKO
NEWBY T.J. *see* MORGAN K.L.
NEWBY T.J., STOKES C.R. & BOURNE F.J. Effects of feeding bacterial lipopolysaccharide and dextran sulphate on the development of oral tolerance to contact sensitizing agents, 617
NOMOTO K., *see* SAKEMI T.
NOMOTO K., *see* SHIMAMOTO Y.
NOMOTO K., YOSHIDA S. & HIMENO K. Comparability of delayed hypersensitivity in various rodents. II. Jones–Mote type hypersensitivity in guinea-pigs immunized with sheep erythrocytes and its modification by cyclophosphamide or BCG pre-treatment, 83

OGDEN B.E. & HILL H.R. Histamine regulates lymphocyte mitogenic responses through activation of specific H_1 and H_2 histamine receptors, 107
OKUDA T. & TACHIBANA T. Complement receptors on Raji cells. The presence of a new type of C3 receptor, 159
OLDROYD R.G., *see* LACHMANN P.J.
OSAWA T., *see* SAWADA J.
OSHIMI K., *see* KANO S.
OTTAWAY C.A., MANSON-SMITH D.F., BRUCE R.G. & PARROTT DELPHINE M.V. Regional blood flow and the localization of lymphoblasts in the small intestne of the mouse. II. The effect of a primary enteric infection with *Trichinella spiralis*, 963
OTTAWAY C.A. & PARROTT DELPHINE M.V. Regional blood flow and the localization of lymphoblasts in the small intestine of the mouse. I. Examination of normal small intestine, 955
OUDENAREN A. VAN, *see* HAALJMAN J.J.

PANDEY R., *see* KAUSHIK A.K.
PAPAZOGLOU S., *see* DE CEULAER C.
PARISH C.R., *see* HOGG NANCY
PARROTT DELPHINE M.V., *see* OTTAWAY C.A.
PARROTT DELPHINE, *see* ROCHA BENEDITA
PASSWELL J.H., COLTEN H.R., SCHNEEBERGER E.E., MAROM Z. & MERLER E. Modulation of human monocyte functions by Fc fragments of IgG: a comparison to other monocyte 'activators', 217
PAYNE A.W.R., *see* HALL J.G.
PEPYS M.B., *see* BALTZ MARILYN L.
PERDIGÓN GABRIELA, *see* MARGNI R.A.
PHONDKE G.P., *see* SELVAN EMILY
PLAYFAIR J.H.L., *see* DOCKRELL HAZEL M.
PLAYFAIR J.H.L., *see* BRITTLE M.P.
PLUMB D., *see* EREMIN O.
POPHAM ANGELA M., *see* DRESSER D.W.
POPHAM ANGELA M. & DRESSER D.W. Rheumatoid factors in mice: non-specific activators of heterophile rheumatoid factor production, 579
PTAK W., *see* MARCINKIEWICZ J.

RENSHAW H.W. & GILMORE R.J. Alternative and classical complement pathway activity in sera from colostrum-fed and colostrum-deprived neonatal pigs, 203
RETZEL E.F., *see* BUMOL T.F.
RHODES JOAN M. & LIND INGA Uptake and degradation of staphylococcal protein A–IgG and HSA–IgG complexes by peritoneal macrophages from rabbits, guinea-pigs and mice, 75
RICCI M., *see* ROMAGNANI S.
RICHTER M., *see* CORTENS M.
RIJKERS G.T., FREDERIX-WOLTERS ELIZABETH M.H. & MUISWINKEL W.B. VAN The immune system of cyprinid fish. Kinetics and temperature dependence of antibody-producing cells in carp (*Cyprinus carpio*), 91
ROCHA BENEDITA, FREITAS A.A. & PARROTT DELPHINE Characterization of rat spleen cell populations. II. Activation by antigens and allogeneic cells, 825
RODER, J.C., Target-effector interaction in the natural killer (NK) cell system VI. The influence of age and genotype on NK binding characteristics, 483
ROMAGNANI S., DEL PRETE G.F., MAGGI E., FALAGIANI P. & RICCI M. T-cell independence of immunoglobulin synthesis by human peripheral blood lymphocytes stimulated with SpA-containing staphylococci, 921
RON Y., *see* DE BAETSELIER P.
ROOK G.A.W., *see* BAHR G.M.
RUSSELL M.W., *see* SCULLY C.M.
RYFFEL B., DONATSCH P., GÖTZ U. & TSCHOPP MARIANNE. Cyclosporin receptor on mouse lymphocytes, 913
RYTER ANTOINETTE, *see* BENOLIEL ANNE MARIE
RYVAR RUTH, *see* BEARD. H.

SAINT-REMY J.-M.R. & COLE P. Interactions of chrysotile asbestos fibres with the complement system, 431
SAITO KAZUKO, NATSUNO T. & MITSUHASHI S. Cell participation in immune response by immune ribonucleic acid. I. The role of T lymphocytes in immune response by immune RNA against T-dependent antigen, 937
SAITO KAZUKO, NATSUNO T. & MITSUHASHI S. Cell participation in immune response by immune ribonucleic acid. II. Independence of T lymphocytes in immune response against T-independent antigen, 947
SAKEMI T., KUROIWA A. & NOMOTO K. Effect of carrageenan on the induction of cell-mediated cytotoxic responses *in vivo*, 297
SALAMAN M.R., *see* SARGENT I.L.
SARGENT I.L. & SALAMAN M.R. Effects of human transfer

factor on the migration of guinea-pig macrophages: is there an antigen-specific activity? 227
SAWADA J. & OSAWA T. Characterization of lectin-induced cellular cytotoxicity mediated by mouse spleen cells and the role of lymphotoxin, 525
SAWADA T. & BITO Y. A B-cell subpopulation relatively insusceptible to the cytotoxic effect of anti-bursa cell serum and partially defective in immune function, 331
SCHELLENBERG R.R. Enhanced phospholipid metabolism in rat mast cells stimulated to release histamine, 123
SCHEPER R.J., *see* BERG W.B. VAN DEN
SCHNEEBERGER E.E., *see* PASSWELL J.H.
SCHONFELD S.E. & HERLES SUSAN *In vitro* and *in vivo* autoimmune response to enamel matrix proteins, 475
SCHWARTZ M., *see* LIFSHITZ RUTH
SCULLY C.M., RUSSELL M.W. & LEHNER T. Specificity of opsonizing antibodies to antigens of *Streptococcus mutans*, 467.
SELL K.W., *see* HARE J.A.
SELVAN EMILY, JOSHI D.S. & PHONDKE G.P. A subset of murine T lymphocytes forming rosettes with rabbit erythrocytes, 415
SHER R., *see* WADEE A.A.
SHIMAMOTO Y., TANIGUCHI K., KUBO C. & NOMOTO K. Differences in thymus-dependency among various T-cell functions, 167
SHOHAM J., COHEN MIRIAM, CHANDALI YARDENA & AVNI A. Thymic hormonal activity on human peripheral blood lymphocytes, *in vitro*. I. Reciprocal effect on T and B rosette formation, 353
SHORT HELEN B., *see* KLIPSTEIN F.A.
SHUKLA MANOHAR I., *see* CHATURVEDI U.C.
SIMIĆ M.M., *see* LUKIĆ M.L.
SKLAR SUZI, *see* CORTENS M.
SLINGERLAND-TEUNISSEN JOKE, *see* HAAIJMAN J.J.
SMILLIE D.M., *see* HENDERSON D.C.
SÖDERBERG A., *see* AARDEN L.
SOUTHERN M.A., *see* BROOM B.C.
SOUZA J.B. DE, *see* DOCKRELL HAZEL M.
SPIK GENEVIÈVE, *see* STEPHENS SUSAN
STANFORD J.L., *see* BAHR G.M.
STANWORTH D.R., *see* BATCHELOR K.W.
STIEBER ANNA, *see* GOUD B.
STEENSGAARD J., *see* JOHANSEN A.S.
STEIN H., *see* GERDES J.
STEPHENS SUSAN, DOLBY JEAN M., MONTREUIL J. & SPIK GENEVIÈVE. Differences in inhibition of the growth of commensal and enteropathogenic strains of *Escherichia coli* by lactotransferrin and secretory immunoglobulin A isolated from human milk, 597
STEVENSON F.K., MORRIS D. & STEVENSON G.T. Immunoglobulin produced by guinea-pig leukaemic B lynphocytes: its source and use as a monitor of tumour load, 313
STEVENSON G.T., *see* STEVENSON F.K.
STEWARD M.W., *see* DEVEY MADELEINE E.
STOKES C.R., *see* NEWBY T.J.
SUNDQVIST K.-G. & WANGER LENA. Anchorage and lymphocyte function. Contact-induced augmentation of T-cell activation, 883
SUYIMA M., *see* KANO S.
SYMONS D.B.A. & CLARKSON CAROL A. Isoelectric heterogeneity of immunoglobulin secreted by LPS-activated neonatal and adult pig lymphocytes *in vitro*, 875

TACHIBANA T., *see* OKUDA T.
TANIGUCHI K., *see* SHIMAMOTO Y.
TAUSSIG M.J. Antigen-specific T-cell factors (review), 759
TRANUM-JENSEN J., *see* BHAKDI S.
TRISSL DOROTHEA, *see* DIAMANTSTEIN T.
TSCHOPP MARIANNA, *see* RYFFEL B.

VAECK M., *see* DE BAETSELIER P.
VEGA J., *see* MARTIN MARIA J.
VALONE F.H. & GOETZL E.J. Structural requirements for the activity of an immunologically generated lipid chemotactic factor, 517
VESOLE D.H. & MOORE G.E. The induction of human peripheral blood lymphoid colonies by conditioned media from human tumour cell lines, 723
VIKSLER V. KH., *see* LAVROVSKY V.A.

WADEE A.A. & SHER R. The effects of a soluble factor released by sensitized mononuclear cells incubated with *S. haematobium* ova on eosinophil migration, 989
WANGER LENA, *see* SUNDQVIST K.-G.
WEBB P.J., *see* BROOKS C.G.
WEINRIEB I.J., *see* MACDERMOTT R.P.
WHALEY K., *see* DE CEULAER C.
WHITE R.G., *see* HENDERSON D.C.
WILKINSON P.C. Effects of human IgG on locomotion of human neutrophils related to IgG binding of a hydrophobic probe, 457
WILLIAMS A.J., HASTINGS M.J.G., EASMON C.S.F. & COLE P.J. Factors affecting the *in vitro* assessment of opsonization: a study of the kinetics of opsonization using the technique of phagocytic chemiluminescence, 903
WILLIAMS A.J., *see* EASMON C.S.F.
WRIGHT B., *see* HOWARD J.C.
WRIGHT B.W., *see* LACHMANN P.J.

YEFENOV E. *In vitro* generation of cytotoxic lymphocytes against radiation and radiation leukaemia virus induced tumours. II. A radiation-induced thymoma generates cytotoxic response in syngeneic but not in allogeneic lymphocytes, 99
YOSHIDA S., *see* NOMOTO K.

ZAFAR M.N., *see* FOA R.
ZANDERS E.D., LAMB J.R., KONTIAINEN SIRKKA & LAHNER T. Partial characterization of murine and monkey helper factor to a streptococcal antigen, 587

Subject Index

Albumin, human serum, kinetics of the decline phase of the primary response, in the bird, 449
Alloantibodies, isolation of six monoclonal, against rat histocompatability antigens: clonal competition, 131
Alloantigen-primed cells, limiting dilution analysis of the suppressive effect mediated by, 399
Allogeneic cells, activation by antigens and, characterization of rat spleen cell populations. II. 825
Alternative complement pathway, a low molecular weight inhibitor of the, I. Its isolation from human urine and the reaction mechanism, 789
Alveolar macrophages. V. Comparative studies on the antigen presentation activity of guinea-pig and rat alveolar macrophages, 361
Amyloid P-component in mice injected with casein: identification in amyloid deposits and in the cytoplasm of hepatocytes, 59
Antibodies, immunobiological behaviour of rabbit precipitating and non-precipitating (co-precipitating), 681
Antibodies, specificity of opsonizing, to antigens of *Streptococcus mutans*, 467
Antibodies, three rat monoclonal, to human C3, 503
Antibody to protein antigens, the induction of chronic antigen–antibody complex disease in selectively bred mice producing either high or low affinity, 303
Antibody-dependent and spontaneous cell-mediated cytotoxicity, are mediated by different processes or cell types, further evidence that, 439
Antibody-dependent cellular cytotoxicity, study of the effect of single or multiple doses of BCG on, 687
Antibody-independent complement activation by a subpopulation of mitogen-induced human T lymphoblasts, 535
Antibody-producing cells in carp (*Cyprinus carpio*), kinetics and temperature dependence of, the immune system of cyprinid fish, 91
Antibody response to SRBC in rats, accessory cell function in immune responses *in vivo*: enhancing effect of radioresistant spleen cells on, 323
Antibody synthesis, IgE, in experimental animals and man, recent studies on the regulation of, 1
Anti-bursa cell serum and partially defective in immune function, a B-cell subpopulation relatively insusceptible to the cytotoxic effect of, 331
Antigen–antibody complex disease, the induction of chronic, in selectively bred mice producing either high or low affinity antibody to protein antigens, 303
Antigen binding and endocytosis by immunocytes, a morphological and functional study on, 833
Antigen-induced proliferation assay for rabbit T lymphocytes. I. Characteristics of the response, 997
Antigen-specific splenic suppressor cells, auto, lack of, in mice with an X-linked B-lymphocyte defect, 815
Antigen-specific T-cell factors (review), 759
Antigens and allogeneic cells, activation by, characterization of rat spleen cell populations. II, 825
Antigens, murine thymus-independent and thymus-dependent, *in vitro* and *in vivo* response of lymphoid cells from LHC hamsters to, 705
Antigens, surface, of the murine cytostatic peritoneal macrophage, 187
Asbestor fibres, interactions of chrysotile, with the complement system, 431
Autoantibodies in (NZB × NZW)F_1 mice, sex differences in the regulation of experimentally induced, 819
Autoimmune response to enamel matrix proteins, *in vitro* and *in vivo*, 475

B-cell subpopulation, a, relatively insusceptible to the cytotoxic effect of anti-bursa cell serum and partially defective in immune function, 331
BCG, study of the effect of single or multiple doses of, on antibody-dependent cellular cytotoxicity, 687
B-lymphocyte defect, X-linked, lack of auto antigen-specific splenic suppressor cells in mice with an, 815
B lymphocytes, immunoglobulin produced by guinea-pig leukaemic, its source and use as a monitor of tumour load, 313
Blood lymphoid colonies, the induction of human peripheral, by conditioned media from human tumour cell lines, 723
Buffalopox virus, cellular and humoral immune responses to, in experimentally infected mice and rabbits, 153

C1, the relationship between the binding ability and the rate of activation of the complement component, 179
C3, three rat monoclonal antibodies to human, 503
C3, isolation and characterization of rat complement factor B and its interaction with cell-bound human, 849
C3 receptor, complement receptors on Raji cells. The presence of a new type of, 159
C3b receptors, physicochemical characterization of, isolated from human erythrocytes by immunoprecipitation, 929
C5b-9 complement complex, re-incorporation of the terminal, into lipid bilayers: formation and stability of reconstituted liposomes, 737
Carp (*Cyprinus carpio*), the immune system of cyprinid fish. Kinetics and temperature dependence of antibody-producing cells in, 91
Carrageenan, effect of, on the induction of cell-mediated cytotoxic responses *in vivo*, 297
Casein, amyloid P-component in mice injected with: identification in amyloid deposits and in the cytoplasm of hepatocytes, 59
Chemotactic factor, structural requirements for the activity of an immunologically generated lipid, 517

Chrysotile asbestos fibres, interactions of, with the complement system,
Collagen types and proteoglycan in pig intervertebral discs, immunochemical localization of, 491
Colostrum-fed and colostrum-deprived neonatal pigs, alternative and classical complement pathway activity in sera from, 203
Complement activation by a subpopulation of mitogen-induced human T lymphoblasts, antibody-independent, 535
Complement component CI, the relationship between the binding ability and the rate of activation of the, 179
Complement components, increased biosynthesis of, by cultured monocytes, synovial fluid macrophages and synovial membrane cells from patients with rheumatoid arthritis, 37
Complement factor B, isolation and characterization of rat, and its interaction with cell-bound human C3, 849
Complement pathway, alternative, a low molecular weight inhibitor of the, I. Its isolation from human urine and the reaction mechanism, 789
Complement, pathway alternative, in bovine serum, variation in the susceptibility of bovine mycoplasmas, 561
Complement pathway activity, alternative and classical, in sera from colostrum-fed and colostrum-deprived neonatal pigs, 203
Complement receptors on Raji cells. The presence of a new type of C3 receptor, 159
Complement system, interactions of chrysotile asbestos fibres with the, 431
Complexes with different cell populations, 21
Con-A induced T-cell proliferation, influence of various macrophage populations on, 379
Contact sensitivity, macrophage suppressor factorin. Mechanisms of its release and action, 211
Contact sensitizing agents, effects of feeding bacterial lipopolysaccharide and dextran sulphate on the development of oral tolerance to, 617
Cyclophosphamide or BCG pre-treatment, Jones–Mote type hypersensitivity in guinea-pigs immunized with sheep erythrocytes and its modification by, comparability of delayed hypersensitivity in various rodents. II, 83
Cyclosporin receptor on mouse lymphocytes, 913
Cyprinid fish, the immune system in, kinetics and temperature dependence of antibody-producing cells in Carp (*Cyprinus carpio*), 91
Cytotoxic agents, the effects of. Properties of peritoneal exudate lymphocytes that mediate tuberculin delayed-type hypersensitivity and anti-tuberculosis immunity. I, 635
Cytotoxic effect of anti-bursa cell serum and partially defective in immune function, a B-cell subpopulation relatively insusceptible to the, 331
Cytotoxic factor, characterization of the, produced in the spleen of dengue virus-infected mice, 387
Cytotoxic reaction towards erythrocytes of various species, the detection and identification of subpopulations of circulating human lymphocytes, monocytes and neutrophils capable of effecting a mitogen-induced cell-mediated, 623
Cytotoxic responses *in vivo*, effect of carrageenan on the induction of cell-mediated, 297
Cytotoxicity, characterization of lectin-induced cellular, mediated by mouse spleen cells and the role of lymphotoxin, 525
Cytotoxicity, further evidence that antibody-dependent and spontaneous cell-mediated, are mediated by different processes or cell types, 439
Cytotoxicity, natural, in the guinea-pig: the natural killer (NK) cell activity of the Kurloff cell, 367
Cytotoxicity, study of the effect of single or multiple doses of BCG on antibody-dependent cellular, 687

Delayed hypersensitivity, comparability of, in various rodents. II. Jones–Mole type hypersensitivity in guinea-pigs immunized with sheep erythrocytes and its modification by cyclophosphamide or BCG pre-treatment, 83
Delayed hypersensitivity reactions, suppression of antibody-mediated accumulation of eosinophils in chronic inflammatory lesions by concomitant, 981
Delayed-type hypersensitivity and anti-tuberculosis immunity, properties of peritoneal exudate lymphocytes that mediate tuberculin, I. The effect of cytotoxic agents, 635
Delayed-type hypersensitivity, properties of peritoneal exudate lymphocytes that mediate tuberculin, II. Velocity sedimentation analysis, 643
Dengue virus-infected mice, characterization of the cytotoxic factor produced in the spleen of, 387
Dental caries, oral immunization with *Streptococcus mutans* in rhesus monkeys and the development of immune responses and, 857
Deoxynucleotidyl transferase expression, terminal, in a Thy-1 alloantigen variant lymphoma cell line, 799
Dextran sulphate, effects of feeding bacterial lipopolysaccharide and, on the development of oral tolerance to contact sensitizing agents, 617

ELISA, use of the, to screen for anti-thymocyte and anti-β_2 microglobulin antibodies in leprosy and SLE, 865
Enamel matrix proteins, *in vitro* and *in vivo* autoimmune response to, 475
Endocytosis by immunocytes, a morphological and functional study on antigen binding and, 833
Entamoeba histolytica extracts, mitogenicity of, for murine lymphocytes, 347
Enterotoxins, immunological cross-reactivity of heat-labile, produced by enterotoxigenic and enteropathogenic strains of *Escherichia coli*, 115
Eosinophil migration, the effects of a soluble factor released by sensitized mononuclear cells incubated with *S. haematobium* ova on, 989
Eosinophils, suppression of antibody-mediated accumulation of, in chronic inflammatory lesions by concomitant delayed hypersensitivity reactions, 918
Erythrocytes, human, physiochemical characterization of C3b receptors isolated from, by immunoprecipitation, 929
Erythrocytes, a subset of murine T lymphocytes forming rosettes with rabbit, 415
Erythrocytes of various species, the detection and identification of subpopulations of circulating human lymphocytes, monocytes and neutrophils capable of effecting a mitogen-induced cell-mediated cytotoxic reaction towards, 623
Escherischia coli, immunoloical cross-reactivity of heat-labile enterotoxins produced by enterotoxigenic and enteropathogenic strains of, 107
Escherichia coli, differences in inhibition of the growth of commensal and enteropathogenic strains of, by lactotrans-

ferrin and secretory immunoglobulin A isolated from human milk, 597

Fc fragments of IgG, modulation of human monocyte functions by, a comparison to other monocyte 'activators', 217

Genes, specificity of, controlling immune responsiveness to (T,G)-A–L and (Phe,G)-A–L, 339
Glutaraldehyde fixation, the effect of, on the immunogenicity of allogeneic lymphoid and tumour cells, 715

Helper factor to a streptococcal antigen, partial characterization of murine and monkey, 587
Hepatocytes, amyloid P-component in mice injected with casein: identification in amyloid deposits and in the cytoplasm of, 59
Histamine, enhanced phospholipid metabolism in rat mast cells stimulated to release, 123
Histamine receptors, histamine regulates lymphocyte mitogenic responses through activation of specific H_1 and H_2, 107
Histamine regulates lymphocyte mitogenic responses through activation of specific H_1 and H_2 histamine receptors, 107
Histamine-releasing properties, a comparison of the, of rat pleural and peritoneal mast cells, 271
Histocompatability antigens, isolation of six monoclonal alloantibodies against rat, clonal competition, 131
Human serum albumin in the bird, kinetics of the decline phase of the primary response, 449
Human transfer factor, effect of, on the migration of guinea-pig macrophages: is there an antigen-specific activity? 227
Hydrophobic probe, effects of human IgG on locomotion of human neutrophils related to IgG binding of a, 457
Hypersensitivity, and anti-tuberculosis immunity. II. Properties of peritoneal exudate lymphocytes that mediate tuberculin delayed-type, velocity sedimentation analysis, 643
Hypersensitivity, and anti-tuberculosis immunity. I. Properties of peritoneal exudate lymphocytes that mediate tuberculin delayed-type, the effects of cytotoxic agents, 635
Hypersensitivity, delayed reactions, suppression of antibody-mediated accumulation of easinophils in chronic inflammatory lesions by concomitant, 981
Hypersensitivity, Jones–Mote type, in guinea-pigs immunized with sheep erythrocytes and its modification by cyclophosphamide or BCG pre-treatment, comparability of delayed hypersensitivity in various rodents. II, 83

Ig levels, kinetics of recovery of serum, and of cytoplasmic Ig positive cells in various lymphoid organs of nude mice after thymus transplantation, 279
Ig, the three major classes of, *in vitro* response of human peripheral blood lymphocytes to protein A DNA synthesis and generation of cells synthesizing, 891
Ig positive cells in various lymphoid organs of nude mice after thymus transplantation, kinetics of recovery of serum Ig levels and of cytoplasmic, 279
IgE antibody synthesis in experimental animals and man, recent studies on the regulation of, 1
IgGa, binding characteristics of aggregated, to rat basophilic leukaemia (RBL) rat cells and rat mast cells, 605
IgG complexes, uptake and degradation of staphylococcal protein A–IgG and HSA–, by peritoneal macrophages from rabbits, guinea-pigs and mice, 75
IgG, effects of human, on locomotion of human neutrophils related to IgG binding of a hydrophobic probe, 457
IgG, Fc fragments of, modulation of human monocyte functions by, a comparison to other monocyte 'activators', 217
IgG immune complexes, solubility properties of, comparison between the effects of low molecular weight solvents and polyethylene glycols, 695
Immunochemical localization of collagen types and proteoglycan in pig intervertebral discs, 491
Immunocytes, a morphological and functional study on antigen binding and endocytosis by, 833
Immunoglobulin A, secretory, isolated from human milk, differences in inhibition of the growth of commensal and enteropathogenic strains of *Escherichia coli* by lactotransferrin and, 597
Immunoglobulin A from blood to bile, comparative aspects of the transport of, 899
Immunoglobulin class and subclass, the role of T cells in regulating, 743
Immunoglobulin, isoelectric heterogeneity of, secreted by LPS-activated neonatal and adult pig lymphocytes *in vitro*, 875
Immunoglobulin produced by guinea-pig leukaemic B lymphocytes: its source and use as a monitor of tumour load, 313
Immunoglobulin-related molecules, demonstration of, on human peripheral blood T lymphocytes using chicken anti-F $(ab')_2$ and anti-IgM antibodies, 663
Immunoglobulins, quantitation and origin of the, in porcine respiratory tract secretions, 729
Immunoglobulin synthesis, T-cell independence of, by human peripheral blood lymphocytes stimulated with SpA-containing staphylococci, 921
Inhibition of the growth of commensal and enteropathogenic strains of *Escherichia coli* by lactotransferrin and secretory inmmunoglobulin A isolated from human milk, differences in, 597
Inhibitors, non-specific binding by macrophages: existence of different adhesive mechanisms and modulation by metabolic, 547
Inhibitory processes, non-specific, of immunological and mitogenic cellular responses. I. Comparative effect of four suppressive agents, 143
Intestine, regional blood flow and the localization of lymphoblasts in the small intestine of the mouse. II. The effect of primary enteric infection with *Trichinella spiralio*, 963
Intestine, regional blood flow and the localization of lymphoblasts in the small intestine of the mouse. I. Examination of normal small, 955

Killer cell, the nature of the, depends on the type of assay, quantitative studies of natural immunity to solid tumours in rats, 673
Kurloff cell, natural cytotoxicity in the guinea-pig: the natural killer (NK) cell activity of the, 367

Lactotransferrin and secretory immunoglobulin A isolated from human milk, differences in inhibition of the growth of commensal and enteropathogenic strains of *Escherichia coli* by, 597

Lectin-induced cellular cytotoxicity, characterization of, mediated by mouse spleen cells and the role of lymphotoxin, 525
Leprosy and SLE, use of the ELISA to screen for anti-thymocyte and anti-β_2 microglobulin antibodies in, 865
Leukaemia, basophilic rat cells (RBL) and rat mast cells, binding characteristics of aggregated IgGa to, 605
Leukaemic B lymphocytes, immunoglobulin produced by guinea-pig, its source and use as a monitor of tumour load, 313
Lidocaine, the effect of, on the processing of soluble immune aggregates and immune complexes by peritoneal macrophages, 237
Lipid chemotactic factor, structural requirements for the activity of an immunologically generated, 517
Lipopolysaccharide, effects of feeding bacterial, and dextran sulphate on the development of oral tolerance to contact sensitizing agents, 617
Lipopolysaccharide (LPS), effect of, on the thymocyte response to PHA: strain difference, 393
Liposomes, re-incorporation of the terminal C5b-9 complement complex into lipid bilayers: formation and stability of reconstituted, 737
Liver, the role of the, in immunity to blood-stage murine malaria, 421
Lymphoblasts, regional blood flow and the localization of, in the small intestine of the mouse. I. Examination of normal small intestine, 955
Lymphoblasts, regional blood flow and the localization of, in the small intestine of the mouse. II. The effect of a primary enteric infection with *Trichinella spiralis*, 963
Lymphocyte function, anchorage and, contact-induced augmentation of T-cell activation, 883
Lymphocyte mitogenic responses, histamine regulates, through activation of specific H_1 and H_2 histamine receptors, 107
Lymphocytes, activation of secondary cytotoxic, by cell-free factors from *I*-region-primed and *D*-region-primed lymphocytes, 653
Lymphocytes, cyclosporin receptor on mouse, 913
Lymphocytes, cytotoxic, *in vitro* generation of, against radiation and radiation leukaemia virus induced tumours. II. A radiation-induced thymoma generates cytotoxic response in syngeneic but not in allogeneic lymphocytes, 99
Lymphocytes, human peripheral blood T, demonstration of immunoglobulin-related molecules on, using chicken anti-$F(ab')_2$ and anti-IgM antibodies, 663
Lymphocytes *in vitro*, isoelectric heterogeneity of immunoglobulin secreted by LPS-activated neonatal and adult pig, 875
Lymphocytes, *in vitro* response of human peripheral blood, to protein A DNA synthesis and generation of cells synthesizing the three major classes of Ig, 891
Lymphocytes, mitogenicity of *Entamoeba histolytica* extracts for murine, 347
Lymphocytes, monocytes and neutrophils capable of effecting a mitogen-induced cell-mediated cytotoxic reaction towards erythrocytes of various species, the detection and identification of subpopulations of circulating human, 623
Lymphocytes, properties of peritoneal exudate, that mediate tuberculin delayed-type hypersensitivity and anti-tuberculosis immunity. II. Velocity sedimentation analysis, 643
Lymphocytes, properties of peritoneal exudate, that mediate tuberculin delayed-type hypersensitivity and anti-tuberculosis immunity. I. The effect of cytotoxic agents, 635
Lymphocytes stimulated with SpA-containing staphylococci, T-cell independence of immunoglobulin synthesis by human peripheral, 921
Lymphocytes, thymic hormonal activity on human peripheral blood, *in vitro*. I. Reciprocal effect on T and B rosette formation, 353
Lymphoid cells from LHC hamsters, *in vitro* and *in vivo* response of, to murine thymus-independent and thymus-dependent antigens, 705
Lymphoid colonies, the induction of human peripheral blood, by conditioned media from human tumour cell lines, 723
Lymphoid and tumour cells, the effect of glutaraldehyde fixation on the immunogenicty of allogeneic, 715
Lymphoid organs of nude mice after thymus transplantation, kinetics of recovery of serum Ig levels and of cytoplasmic Ig positive cells in various, 279
Lymphoma cell line, terminal deoxynucleotidyl transferase expression in a Thy-1 alloantigen variant, 663
Lymphotoxin, characterization of lectin-induced cellular cytotoxicity mediated by mouse spleen cells and the role of, 525

Macrophage populations, influence of various, on Con-A induced T-cell proliferation, 379
Macrophage, surface antigens of the murine cytostatic peritoneal, 187
Macrophage suppressor factor in contact sensitivity. Mechanisms of its release and action, 211
Macrophages, alveolar, V. Comparative studies on the antigen presentation activity of guinea-pig and rat alveolar macrophages, 361
Macrophages, effects of human transfer factor on the migration of guinea-pig, is there an antigen-specific activity? 227
Macrophages, the effect of lidocaine on the processing of soluble immune aggregates and immune complexes by peritoneal, 237
Macrophages, non-specific binding by, existence of different adhesive mechanisms and modulation by metabolic inhibitors, 547
Macrophages, synovial fluid, monocytes and synovial membrane cells from patients with rheumatoid arthritis, increased biosynthesis of complement components by cultured, 37
Malaria, the role of the liver in immunity to blood-stage murine, 421
Mast cells, binding characteristics of aggregated IgGa to rat basophilic leukaemia (RBL) cells and rat, 605
Mast cells, growth and differentiation *in vitro* of, from mesenteric lymph nodes of *Nippostrongylus brasiliensis*-infected rats, 195
Mast cells, a comparison of the histamine-releasing properties of rat pleural and peritoneal, 271
Mast cells, rat, enhanced phospholipid metabolism in , stimulated to release histamine, 123
Meynell, superinfecting phage technique of, the *in vivo* division and death rates of Salmonella typhimurium in

the spleens of naturally resistant and susceptible mice measured by the, 973
Mitogenicity of *Entamoeba histolytica* extracts for murine lymphocytes, 347
Mitogen-induced human T lymphoblasts, antibody-independent complement activation by a subpopulation of, 535
Monocyte 'activators', a comparison to other, modulation of human monocyte functions by Fc fragments of IgG, 217
Monocyte functions, modulation of human, by Fc fragments of IgG: a comparison to other monocyte 'activators', 217
Monocytes, bovine peripheral blood, the isolation, long-term cultivation and characterization of, 807
Monocytes, lymphocytes and neutrophils capable of effecting a mitogen-induced cell-mediated cytotoxic reaction towards erythrocytes of various species, the detection and identification of subpopulations of circulating human, 623
Monocytes, synovial fluid macrophages and synovial membrane cells from patients with rheumatoid arthritis, increased biosynthesis of complement components by cultured, 37
Mucosal immunology (review), 249
Mycoplasmas, variation in the susceptibility of bovine, to killing by the alternative complement pathway in bovine serum, 561

Natural killer activity of Wistar rat spleen cells: blocking effect of homologous sera, 541
Natural killer binding characteristics, the influence of age and genotype on, target-effector interaction in the natural killer (NK) cell system VI, 483
Neutrophils, lymphocytes and monocytes capable of effecting a mitogen-induced cell-mediated cytotoxic reaction towards erythrocytes of various species, the detection and identification of subpopulations of circulating human, 623
Neutrophils, effects of human IgG on locomotion of human, related to IgG binding of a hydrophobic probe, 457
Nippostrongylus brasiliensis-infected rats, growth and differentiation *in vitro* of mast cells from mesenteric lymph nodes of, 195

Opsonic and phagocytic function, the measurement of, by Luminol-dependent chemiluminescence, 67
Opsonization, factors affecting the *in vitro* assessment of, a study of the kinetics of opsonization using the technique of phagocytic chemiluminescence, 903

PHA, effect of lipopolysaccharide (LPS) on the thymocyte response to, strain difference, 393
Phagocytic chemiluminescence, factors affecting the *in vitro* assessment of opsonization: a study of the kinetics of opsonization using the technique of, 903
Phagocytic function, the measurement of opsonic and, by Luminol-dependent chemiluminescence, 67
(Phe,G)-A–L, specificity of genes controlling immune responsiveness to (T,G)-A–L and, 339
Phospolipid metabolism, enhanced, in rat mast cells stimulated to release histomine, 123
Pneumococcal polysaccharides, maternal–foetal interaction, antibody formation, and metabolic response in mice immunized with, 45
Polysaccharides, pneumococcal, maternal–foetal interaction, antibody formation, and metabolic response in mice immunized with, 45
Protein A DNA synthesis and generation of cells synthesizing the three major classes of Ig, *in vitro* response of human peripheral blood lymphocytes to, 891
Proteoglycan and collagen types in pig intervertebral discs, immunochemical localization of, 491

RNase-sensitive thymocytic extract, conversion of immune response patterns from high to low and low to high by an, 25
Raji cells, complement receptors on, the presence of a new type of C3 receptor, 159
Respiratory tract secretions, quantitation and origin of the immunoglobulins in porcine, 729
Rheumatoid arthritis, increased biosynthesis of complement components by cultured monocytes, synovial fluid macrophages and synovial membrane cells from patients with, 37
Rheumatoid factors in mice: plaque assay for homophile and heterophile rheumatoid factors, 569
Rheumatoid factors in mice: non-specific activators of heterophile rheumatoid factor production, 579
Ribonucleic acid, cell participation in immune response by immune, I. The role of T lymphocytes in immune response by immune RNA against T-dependent antigen, 937
Ribonucleic acid, cell participation in immune response by immune, II. Independence of T lymphocytes in immune response against T-independent antigen, 947
Rosette formation, thymic hormonal activity on human peripheral blood lymphocytes, in vitro: I. Reciprocal effect on T and B, 353
Rosettes with rabbit erythrocytes, a subset of murine T lymphocytes forming, 415

Salmonella typhimurium, the *in vivo* division and death rates of, in the spleens of naturally resistant and susceptible mice measured by the superinfecting phage technique of Meynell, 973
Sera, natural killer activity of Wistar rat spleen cells: blocking effect of homologous, 541
S. haematobium ova on eosinophil migration, the effects of a soluble factor released by sensitized mononuclear cells incubated with, 989
SLE, use of the ELISA to screen for anti-thymocyte and anti-β_2 microglobulin antibodies in leprosy and, 865
Spleen cell populations, characterization of rat, II. Activation by antigens and allogeneic cells, 825
Spleen cells, accessory cell function in immune responses *in vivo*: enhancing effect of radioresistant, on antibody response to SRBC in rats, 323
Spleen cells, Wistar rat, blocking effect of homologous sera on natural killer activity of, 541
Splenic suppressor cells, lack of auto antigen-specific, in mice with an X-linked B-lymphocyte defect, 815
Staphylococcal protein A-IgG and HSA, uptake and degradation of, by peritoneal macrophages from rabbits, guinea-pigs and mice, IgG complexes, 75
Staphylococci, SpA-containing, T-cell independence of immunoglobulin synthesis by human peripheral blood lymphocytes stimulated with, 921
Streptococcal antigen, partial characterization of murine and monkey helper factor to a, 587

Streptococcus mutans in rhesus monkeys, oral immunization with, and the development of immune responses and dental caries, 857
Streptococcus mutans, specificity of opsonizing antibodies to antigens of, 467
Suppression, ratio-dominance model of, an analysis by limiting dilution, 407
Suppressive agents, comparative effect of four, non-specific inhibitory processes of immunological and mitogenic cellular responses. I, 143
Suppressive effect mediated by alloantigen-primed cells, limiting dilution analysis of the, 399

(T,G)-A–L and (Phe,G)-A–L, specificity of genes controlling immune responsiveness to, 339
Target-effector interaction in the natural killer (NK) cell system. VI. The influence of age and genotype on NK binding characteristics, 483
T-cell activation, anchorage and lymphocyte function. Contact-induced augmentation of, 883
T-cell factors, antigen-specific (review), 759
T-cell functions, differences in thymus-dependency among various, 167
T-cell independence of immunoglobulin synthesis by human peripheral blood lymphocytes stimulated with Sp-A containing staphylococci, 921
T-cell proliferation, influence of various macrophage populations on Con-A induced, 376
T-cells, the role of, in regulating immunoglobulin class and subclass, 743
T lymphoblasts, human antibody-independent complement activation by a subpopulation of mitogen-induced, 535
T lymphocytes, antigen-induced proliferation assay for rabbit, I. Characteristics of the response, 997
T-lymphocyte colony formation, inhibition of, by inhibitors of mitochondrial protein synthesis, 55
T lymphocytes, a subset of murine, forming rosettes with rabbit erythrocytes, 415
T lymphocytes in immune response against T-independent antigen, cell participation in immune response by immune ribonucleic acid. II. Independence of, 947
T lymphocytes in immune response by immune RNA against T-dependent antigen, cell participation in immune response by immune ribonucleic acid. I. The role of, 937
Thymic hormonal activity on human peripheral blood lymphocytes, *in vitro*. I. Reciprocal effect on T and B rosette formation, 353
Thymocyte extract, conversion of immune response patterns from high to low and low to high by an RNase-sensitive, 25
Thymocyte response to PHA, effect of lipopolysaccharide (LPS) on the, strain difference, 393
Thymoma, a radiation-induced, generates cytotoxic response in syngeneic but not in allogeneic lymphocytes, *in vitro* generation of cytotoxic lymphocytes against radiation and radiation leukaemia virus induced tumours. II, 99
Thymus-dependency, differences in, among various T-cell functions, 167
Trichinella spiralis, regional blood flow and the localization of lymphoblasts in the small intestine of the mouse. II. The effect of a primary enteric infection with, 963
Tumour cell lines, human, the induction of human peripheral blood lymphoid colonies by conditioned media from, 723
Tumour-inhibiting factor (TIF), presence of a, in sera from normal but not tumour-bearing mice, 289
Tumours in rats, quantitative studies of natural immunity to solid, the nature of the killer cell depends on the type of assay, 673

Velocity sedimentation analysis. Properties of peritoneal exudate lymphocytes that mediate tuberculin delayed-type hypersensitivity and anti-tuberculosis. II, 643

Wistar rat spleen cells: blocking effect of homologous sera on natural killer activity of, 541